

Effects of Pruning Intensity on Grape Yield and Quality of Erciř Grape Cultivar (*Vitis vinifera* L.)

Cüneyt UYAK^{1*} Adnan DOĐAN¹ Sinem DELİKANLIOĐLU¹

¹Department of Horticulture, Faculty of Agriculture, Yüzüncü Yıl University, Van, Turkey

*Corresponding Author:
E-mail:cuneytuyak@gmail.com

Received: January 04, 2016
Accepted: February 19, 2016

Abstract

The present study was conducted to investigate the effects of different pruning intensity treatments on grape yield and quality of Erciř grape cultivar in the year 2014. The gobble-trained grapevines grown on their own roots were subjected to three different pruning intensities during winter pruning [16 buds/grapevine (8 shoots x 2 buds); 24 buds/grapevine (8 shoots x 3 buds); 32 buds/grapevine (8 shoots x 4 buds)]. Compared to low pruning intensity (16 buds/grapevine), high pruning intensity treatments (32 buds/grapevine) increased grapevine yield by 55%, but decreased cluster weight by 17%, cluster width by 25.2%, cluster length by 12.5%, grape width by 10.9%, grape length by 6.7%, 100-grape weight by 22.7% and water soluble dry matter yield by 2.6%.

Keywords: Erciř grape, Pruning intensity, Yield, Quality

INTRODUCTION

Grape is among the mostly cultured fruits worldwide. As it was in several fruit species, there are various cultural practices to improve the yield and quality in grapes. Among these practices, pruning is a significant practice. Pruning is a cultural practice targeting to benefit from vineyards uppermost level through arranging growth and development well balanced with yield and quality. Number of buds to be left over the grapevine in winter pruning (pruning intensity – fruit load) is directly effective on growth, development, yield and quality. The cultivar, purpose of use, rootstock, training system, age and development of grapevine and ecological conditions are the factors with great influences on pruning intensity [1]. A physiological balance to be established and preserved between vigor and yield-quality of vines is directly related to conscious winter and summer pruning. Development and fruit quality is directly influenced by the arrangements made over the yields. Therefore while in winter pruning, grapevines should be loaded with a fruit load (pruning intensity) proportional to their development capacities [1, 2, 3, 4, 5, 6]. Since pruning intensity is influenced by several factors, it is necessary to identify proper pruning intensities for different grape cultivars grown under different conditions.

The present study was conducted to investigate the effects of different pruning intensity treatments on growth, yield and quality of Erciř grape cultivar.

MATERIALS and METHODS

The present study was conducted on Erciř grape cultivar grown in Van region of Turkey in 2014. Experimental vineyard is grown over their own roots, 20 years old and gobble-trained with 2x1.5 m planting spacing. Van region has an altitude of 1725 m. Annual average temperature of the region is 10 °C, annual precipitation is 385.2 mm and annual average relative humidity is 57.3% [7]. Vigor levels of the grapevines were determined based on pruning wood weights in winter pruning. Experiments were conducted on grapevines with close vigor levels. Based on vigor levels, three different pruning densities were applied to grapevines as of low [16 buds/grapevine (2 buds x 8 shoots)], normal

[24 buds/grapevine (3 buds x 8 shoots)] and high [32 buds/grapevine (4 buds x 8 shoots)]. Experiments were conducted in randomized blocks design with three replications with 4 grapevines in each replication. Throughout the experiments, two soil tillage, two flooding irrigation and two fungicide treatments against powdery mildew were performed. Since the shoots exhibit a lateral growth without any external supports, shoot tip pinching was performed just before flowering. Following the treatments, shooting ratio (%), annual branch diameter (cm), annual branch length (cm), number of shoots per grapevine, number of clusters per shoot, grapevine yield (kg), number of clusters per grapevine, average cluster weight (g), cluster width and length (cm), 100-grape weight (g), grape width and length (mm), juice yield (ml/g), water soluble dry matter yield (g), titratable acidity (g/l) and juice pH values were determined. Resultant data were statistically analyzed with SPSS 22.0 software and means were compared with Duncan's multiple range test.

RESULTS and DISCUSSION

Compared to low pruning density, high pruning density treatments decreased shooting ratio by 6.1% (Table 1). Several other researchers also reported decreasing shooting ratios with increasing pruning intensities [8, 9, 10, 11, 12]. High pruning intensity also decreased branch diameter by 13.6% and branch length by 5.1% (Table 1). Similarly, Dardeniz and Kısmalı [13], Çelik [9], Shalan [12], Shoeib [14] and El-Kady et al. [15] reported decreased branch diameter and length with increasing number of buds left in pruning. Compared to low pruning intensity, high pruning intensity decreased number of shoots per grapevine by 87.5% and number of clusters per shoot by 22.4% (Table 1). İlhan [8] indicated that increase in number of buds increased number of shoots per grapevine and decreased number of clusters per shoot in seedless grape cultivar.

Again compared to low pruning intensity, high pruning intensity treatments increased grapevine yield by 55.0% and number of clusters per grapevine by 53.7% and decreased average cluster weight by 17.0% (Table 1). Previous researchers reported increasing grapevine yields and number of clusters per grapevine and decreasing average cluster weights with increasing number of buds left in pruning [8, 9,

12, 16, 17, 18, 19, 20].

It was observed that high pruning intensity decreased cluster width by 25.2%, cluster length by 12.5%, grape width by 10.9%, grape length by 6.7 and 100-grape weight by 22.7% (Table 1). Shalan [12]; Dardeniz and Kismalı [13] reported decreasing cluster dimensions with high pruning intensities; Çelik [9] indicated that pruning intensity was not effective on grape weight, grape volume and dimensions; Ewart et al. [21]; Christensen et al. [22] reported decreasing grape dimensions with decreasing pruning intensities; İlhan [8]; İlgin and Yıldız [19]; Dardeniz and Kismalı [13] indicated that high pruning intensity decreased 100-grape weights.

Compared to low pruning intensity, high pruning intensity treatments decreased juice yield by 2.4%, water soluble dry matter yield by 2.6%, juice pH by 4.1% and increased titratable acidity by 25% (Table 1). Similar results were also

reported by previous researchers [8, 12, 13, 16, 17, 19, 23, 24].

CONCLUSION

It was concluded based on current findings that increase in number of buds increased grapevine yield, number of clusters, number of shoots per grapevine and titratable acidity; decreased grapevine development, grape yield and quality parameters. Decreases in cluster weights resulted from decreases in cluster and grape dimensions and weights and the increase in yield was resulted from increase in number of clusters. Normal pruning intensity (24 buds/grapevine) with medium branch diameters, medium size clusters and grapes, sufficient yield levels was identified as the most proper pruning level for different training systems and vigor levels of vineyard.

Table 1. Effects of different pruning intensities on grapevine development, grape yield and quality

Characteristics	Low (16 buds/grapevine)	Normal (24 buds/grapevine)	High (32 buds/grapevine)	Mean
Shooting ratio (%)	0.95±0.08a	0.96±0.00a	0.90±0.05a	0.94±0.06
Branch diameter (mm)	10.23±2.15a	9.64±1.67b	9.00±1.22c	9.626±1.79*
Branch length (cm)	96.15±16.15a	93.39±19.16a	91.44±20.00a	93.661±18.51
Number of shoots per grapevine (shoot number /grapevine)	15.33±1.23c	22.00±1.21b	28.75±1.66a	22.03±5.72*
Number of clusters per shoot (cluster number /shoot)	1.18±0.17a	1.107±1.10a	0.96±0.08b	1.083±1.50*
Grapevine yield (kg/grapevine)	7.12±1.95b	9.54±2.23a	11.05±2.35a	9.24±2.69*
Number of clusters per grapevine (cluster number / grapevine)	18.00±2.37c	24.33±2.06b	27.67±2.23a	23.33±4.60*
Average cluster weight (g)	353.45±101.83a	335.02±112.55ab	301.94±78.73b	330.1±104.2*
Cluster width (cm)	11.57±2.24a	10.300±1.79b	9.23±1.36c	10.367±2.06*
Cluster length (cm)	19.16±2.85a	18.38±2.40a	17.03±2.21b	18.19±2.64*
100-grape weight (g)	278.47±31.94a	237.90±21.23b	226.94±22.10b	247.77±33.5*
Grape width (mm)	17.58±1.30a	16.44±1.26b	15.85±1.33c	16.62±1.48*
Grape length (mm)	17.08±1.21a	16.28±1.00b	15.99±1.28b	16.45±1.25*
Juice yield (ml/100 g)	73.27±2.66a	73.20±2.96a	71.53±2.90a	72.67±2.89
Water soluble dry matter (%)	17.52±1.10a	17.40±0.34a	17.07±0.75a	17.33±0.80
Titratable acidity (g/l)	5.20±1.19b	6.27±1.04a	6.50±1.36a	5.99±1.31*
Juice pH	3.33±0.042a	3.21±0.05b	3.19±0.08b	3.24±0.08*

The means indicated with different letters in the same row are significantly different ($p<0.05$). * The difference between the means is significant ($p<0.05$).

This study has been supported by Yüzüncü Yıl University BAP Unit.

REFERENCES

- [1] Çelik H, Ağaoğlu YS, Fidan Y, Maraslı B, Söylemezoglu G. 1998. Genel Bağcılık. Sunfidan A.Ş. Mesleki Kitaplar Serisi:1. Ankara. 253.
- [2] Pantic Z. 1975. Contribution to studies on the number of buds left when pruning grapevines. *Nauka v Praksi*, 3(2): 137-142, [Hort. Abstr., 45(8): 5752].
- [3] İler E, Çımrın, T. 1976. Şaraplık üzümelerde çeşit ve sarj faktörünün verim ve kaliteye etkisi üzerine araştırmalar. *Bitki* 3(4): 340-349.
- [4] Nikiforova LT, Martyanova DA, Komarovskii A M, Litunovskii AG. 1981. Rational system of pruning the grapevine cultivar Aliogaté trained on a high stem. *Sadovodstvo*

and *Vinogradarstvo 1 vinodelie Moldavii*, 6: 30-32, [Hort. Abstr., 51(8): 6140].

[5] Jackson DI, Lombard PB. 1993. Environmental and management practices affecting grape composition and wine quality. A review. *Amer. J. Enol. Vitis*, 44: 409-430.

[6] Clingeffer PR. 1989. Update: Minimal pruning of cordon trained vines. *Austr. Grapegrower & Winemaker*, 282: 78-83.

[7] Anonymous 2015. Van Meteoroloji İstasyonları Kayıtları. İl Meteoroloji Müdürlüğü, Van.

[8] İlhan İ. 1988. Çekirdeksiz üzüm Çeşidinde Farklı Sayıda ve Uzunlukta Bayrak Bırakarak Yapılan Değişik Şarjın Asmanın Vegetatif ve Generatif Gelişmesine Etkileri Üzerine Araştırmalar (Doktora Tezi). Ege Üniversitesi, Fen Bilimleri Enstitüsü, Bornova, İzmir.

[9] Çelik G. 1996. Ankara Koşullarında Yetiştirilen Ham-

burg Misketi ve Hafızalı Üzüm Çeşitlerinde Değişik Telli Terbiye Şekillerine Uygulanan Farklı Budama Şiddetinin Gelişme, Verim ve Ürün Kalitesi Üzerine Etkileri (Doktora Tezi). Ankara Üniversitesi, Fen Bilimleri Enstitüsü, Ankara.

[10] Reynolds AG, Edwards CG, Wordle DA., Webster DR, Dever M. 1994. Shoot density affects riesling grapevines. I. Vine performance. *J. Am. Soc. Hortic. Sci.*, 119: 874-880.

[11] Omar AH, Abdel-Kawi A. 2000. Optimal bud load for Thompson Seedless grapevines. *J. Agric. Sci. Mansoura Univ.* 25: 5769-5777.

[12] Shalan AM. 2013. Performance of *Vitis vinifera* cultivar Flame Seedless grapevines under different node load per centimeter square of trunk cross-sectional area. *Asian Journal of Crop Science*, 5(2): 139-152.

[13] Dardeniz A, Kısmalı İ. 2002. Amasya ve Cardinal Üzüm çeşitlerinde farklı ürün yüklerinin üzüm ve çubuk verimi ile kalitesine etkileri üzerine araştırmalar. *Ege Üniversitesi Ziraat Fakültesi Dergisi*, 39 (1): 9-16.

[14] Shoeb MM. 2004. Effect of potassium sulphate and vine load on the growth and yield of Thompson Seedless grapevines with a special reference to the occurrence of cluster tip desiccation problem. *J. Agric. Sci. Mansoura Univ. Egypt*, 29: 4711-4728.

[15] El-Kady MI, Samra NR, El-Shahat AS. 2010. Studies on the effect of potassium and magnesium fertilization as well as vine load on Flame Seedless grape productivity. *J. Plant Prod. Mansoura Univ.*, 1: 1299-1311.

[16] Tokalı S. 1986. Bazı Üzüm Çeşitlerinde Asma Şarjının Üzüm Verimi ve Kalitesine Etkileri Üzerinde Araştırmalar (Yüksek Lisans Tezi). Ege Üniversitesi, Fen Bilimleri Enstitüsü, Bornova, İzmir.

[17] Heil RM. 1998. Effect of Training System and Pruning Severity on The Growth, Yield and Fruit Composition of *Vitis vinifera* L. Cabernet Sauvignon Grapevines. (master thesis). California State University, School of Agricultural Sciences and Technology, Fresno.

[18] Çelik M, Kısmalı İ. 2005. Aşılı ve aşısız Yuvarlak Çekirdeksiz üzüm çeşidi bağlarında farklı şarj uygulamalarının üzüm verimi ve kalitesi ile vegetatif gelişmeye etkileri üzerinde araştırmalar. VI. Bağcılık Sempozyumu. 19-23 Eylül 2005, Tekirdağ. 74-82.

[19] Ilgın C, Yıldız S. 2005. Siyah Kuş üzümü (Black Corinth) çeşidinde farklı göz yükünün üzüm verimi kalitesi üzerine etkisi. VI. Bağcılık Sempozyumu. 19-23 Eylül 2005, Tekirdağ. 398-402.

[20] Pehlivan EC, Uzun Hİ. 2015. Shiraz üzüm çeşidinde salkım seyreltmenin verim ve kalite özellikleri üzerine etkileri. *YYÜ. Tarım Bilimleri Dergisi*, 25 (2): 119-126.

[21] Ewart AJW, Brien CJ, Soderlund R, Smart RE. 1985. The effect of light pruning, irrigation and improved soil management on wine quality of the *Vitis vinifera* cv. Riesling. *Vitis*, 24: 209-217.

[22] Christensen LP, Leavitt MG, Hirschfeld JD, Bianchi LM. 1994. The effects of pruning level and post-bud-break cane adjustment on Thompson Seedless raisin production and quality. *Am. J. Enol. Vitic.*, 45 (2): 141-149.

[23] Fawzi MIF, Shahin MFM, Kandil EA. 2010. Effect of bud load on bud behavior, yield, cluster characteristics and some biochemical contents of the cane of Crimson Seedless grapevines. *J. Am. Sci.*, 6: 187-194.

[24] Rizk MH, Samra NR, El-Kady MI, El-Kenawy M A. 2006. Influence of pruning severity on bud behavior, yield, berry quality and content of total carbohydrates in the

canes of Thompson Seedless grapes under pergola trellis system. *J. Agric. Sci. Mansoura Univ. Egypt*, 31: 901-913.



Figure 1. Leaves of Erciş grape cultivar



Figure 2. Cluster of Erciş grape cultivar



Figure 3. Pruning treatments