

## RESPONSE OF BREAD WHEAT (*TRITICUM AESTIVUM* L) VARIETIES TO SEED SOURCES AND NPSB FERTILIZER RATES FOR GROWTH, YIELD AND SEED QUALITY AT ANCHA SADICHO DISTRICT, SOUTHERN ETHIOPIA

Mulugeta Abebe\*, Yohannes Erkeno

<sup>1</sup>Wachemo University, College of Agricultural Sciences, Department of Plant Science, Hossana, Ethiopia

\*Corresponding Author:

E-mail: [mulewaabe@gmail.com](mailto:mulewaabe@gmail.com)

(Received 23<sup>th</sup> November 2021; accepted 06<sup>th</sup> April 2022)

### ABSTRACT

Wheat is one of the most important cereal crops globally and is a staple food for about one third of the world population. Inappropriate uses of fertilizer rate, unavailability of quality seed for varieties that are high yielding as well as adapted to wide range of agro-ecologies of the area and the use of farm saved poor quality seeds of local varieties are some of the major factors which limit yield of wheat. Therefore, field experiment was conducted during 2020 cropping season to determine the response of bread wheat (*Triticum aestivum* L.) varieties to seed sources and blended fertilizer rates for growth, yield and seed quality in southern Ethiopia. Factorial combinations of four blended NPSB levels (0, 75, 100 and 125 kg ha<sup>-1</sup>), three seed sources (Research, C<sub>2</sub> and C<sub>3</sub>) and three wheat varieties (Hidase, Digalu and Ogolicho) were laid out in randomized complete block design with three replications. Combined analysis of the results revealed that days to 50% heading, days to maturity and germination percentage were significantly affected by the main and their interaction effects of fertilizer rate and seed sources. Grain filling period, plant height and number of kernels were significantly affected by the main effects of fertilizer and their interaction with seed sources whereas spike length, grain yield, 1000 kernels weight, speed of germination, seedling dry weight, seedling length, vigor index i and ii were significantly affected by 3-way interaction. The maximum values of grain yield recorded in interaction of 125 kg ha<sup>-1</sup> NPSB fertilizer rates, seeds obtained from research center and Hidase variety. Therefore, it could be concluded that application of 125 kg blended NPSB fertilizer rate, seeds obtained from research center and Hidase variety were economically profitable in the study area.

**Keywords:** *Blended, grain, productivity, germination, vigour*

### INTRODUCTION

Wheat (*Triticum aestivum* L) is one of the most important cereals cultivated in Ethiopia [1]. It belongs to the family Poaceae, subfamily Pooideae and tribe Triticeae. Wheat is a staple food that provides around 20% of protein and calories consumed worldwide [2]. It has the highest content of protein of all the staple cereals and contains essential minerals, vitamins, and lipids. It is the primary source of protein in developing countries where 1.2 billion people are dependent on wheat for survival [3]. Bread wheat in Ethiopia is used in different forms such as bread, porridge, soup and roasted grain. In addition to the grain, the straw of bread wheat is used for animal feed, hatching roofs and bed decking. Wheat production in 2019 was 764.49 million tons around the globe. It ranks first in the world cereal crops accounting for 30% of all cereal food worldwide. Ethiopia is one of the largest wheat producers in the Sub-Saharan Africa, with yearly estimated production of 4.5 million tons on 1.69 million hectares of land [4]. The total wheat production in the Kambata Tambaro zone is 152,153.6 tons with the productivity of 2.4 tons ha<sup>-1</sup>

in 2018/19 cropping season [4] which is lower than the national average 2.66 tons ha<sup>-1</sup> and the world 2.8 tons ha<sup>-1</sup> [4].

The low yield per hectare is attributed to many factors, among these inappropriate uses of fertilizer rate and unavailability of quality seed for varieties that are high yielding as well as adapted to wide range of agro-ecologies of the country and the use of farm saved poor quality seeds of local varieties are some of the major factors. Poor quality seed results in poor germination and poor crop stands at farm and become a significant factor affecting wheat productivity [5]. The low yield is also allied to the depletion of soil fertility due to continuous nutrient uptake of crops, low fertilizer use and insufficient organic matter application. Nutrient mining due to sub optimal fertilizer use in one hand and unbalanced fertilizer uses on other have favored the emergence of multi nutrient deficiency in Ethiopian soils [6]. On the other hand, the sources of plant nutrients for Ethiopian agriculture over the past five decades have been limited to urea and Diammonium Phosphate fertilizers which contain only nitrogen and phosphorus that may not satisfy the nutrient requirements of crops [7]. Based on the national soil data base, in addition to the macronutrients, due to long year cultivation, some of the micronutrients like sulfur, zinc, boron, and copper are depleted from the soil in the major crop producing area of the country. To avert the situation the Ministry of Agriculture of Ethiopia has been recently introduced a new blended fertilizer (NPSB) containing nitrogen, phosphorous, sulfur and boron with the ratio of (18.9% N, 37.7% P<sub>2</sub>O<sub>5</sub>, 6.95% S, and 0.1% B [7]).

Increasing N fertilizer greatly affected grain yield and quality of wheat protein content [8]. Adequate phosphorus enhances many aspects of plant physiology like photosynthesis, flowering, seed maturity and seed development [9]. Sulfur is an essential nutrient required to build yield and achieve grain quality. Boron is essential for cell division and elongation in meristematic tissues, floral organs and for flower male fertility, pollen tube germination along with it's and seed/fruit formation. Boron deficiency impairs grain setting in wheat, resulting in increased number of open spikelet's and decreased number of grains per spike [10]. Therefore, the use of balanced fertilizers containing both macro and micronutrients, which is based on the site-specific soil fertility assessment, is believed to be one of the solutions for reducing such production constraints. The NPSB fertilizer and seed source and its effects on bread wheat yield components and seed quality have not been studied extensively. Information is lacking on the interaction effects of NPSB fertilizer rate with seed source on bread wheat yield and seed quality. However, limited research has been done to elucidate the response of bread wheat to mineral NPSB fertilizer rates application and seed source in the study area. Therefore, the present study was undertaken with the following objectives: to determine the interaction effects of seed sources and NPSB fertilizer rates on the productivity of bread wheat; to assess the effect of seed sources and NPSB fertilizer rates on the seed quality of bread wheat varieties.

## **MATERIALS AND METHODS**

### ***Description of the experimental site***

The field experiment was conducted at Ancha Sadicho district Farmers Training Centre in Doyogana Woreda, Southern Ethiopia during the main cropping seasons of 2020. The site is located at latitude of 9<sup>0</sup>07' to 9<sup>0</sup>32'N, longitude of 41<sup>0</sup>29' - 41<sup>0</sup> 44'E and an altitude of 1700 - 2800 meters above sea level. Experimental area is classified in to *Dega* and *Weynadega* agro-climatic zones with an average annual temperature of 14.5 °C - 18 °C and 700 - 1100 mm of

annual rain fall [11]. Laboratory experiment was conducted at Wachemo University laboratories as per international seed testing association rules and procedures [12].

### ***Experimental materials***

Three bread wheat varieties namely; Hidase, Digalu and Ogolicho were used as a test crop at experimental site. The seeds were collected from research centre which considered as quality seeds and farmers saved seeds as (C<sub>2</sub> and C<sub>3</sub>) which represent low seed quality. The seeds of the varieties from all sources were collected from the 2018/2019 cropping season harvest. The farmers saved seeds were collected from six farmers each considered three bread wheat varieties in equal proportions as two composite samples for field and laboratory work. Different levels of blended fertilizer in the form of NPSB (18.9% N, 37.7% P<sub>2</sub>O<sub>5</sub>, 6.95% S, and 0.1% B) were used and it was applied at the time of planting.

### ***Treatments and experimental design***

Field experiment with treatments of a factorial combination of four blended NPSB fertilizer levels (0, 75, 100 and 125 kg ha<sup>-1</sup>), three varieties (Hidase, Digalu and Ogolicho) and three seed sources (research, FC<sub>2</sub> and FC<sub>3</sub> seeds) were used. The experiment was laid out in Randomized Complete Block Design (RCBD) in 4 x 3 x 3 factorial arrangements with three replications. Each plot size in the experiment was 2.75 m x 2 m, with 10 rows of 20 cm apart giving a gross plot area of 5.5m<sup>2</sup>. Spacing of 1 m and 0.5 m was maintained between adjacent blocks and plots, respectively. All the required data were collected from the middle eight rows leaving the outer most rows in both sides and some distance at both ends as border at each plot of the experiment on the basis of net harvestable plot area. Laboratory experiment was laid out as Completely Randomized Design with three replications.

### ***Experimental procedure and field management***

Before planting soil samples were taken from experimental site according to the appropriate sampling technique and subjected to soil analysis to determine the chemical and physical properties of the soil. Soil sample was taken from 0 to 30 cm depth of the experimental site by using auger. While taking the sample zigzag method of soil sampling was implemented. A composite sample of 1 kg was prepared from the primary samples through quartering method. After passing through the proper drying and grinding processes composite sample was submitted to soil laboratory for the analysis of physical and chemical properties. Total nitrogen, available phosphorus, available boron, available sulphur, exchangeable potassium, organic matter, soil pH, and Cat ion exchange capacity (CEC) and soil texture are the chemical and physical properties were determined in the laboratory from the sample submitted. The experimental field was prepared before sowing. Seeds were drilled in rows at depth of 4-5 cm. While conducting the experiment all the necessary agronomic and management practices were carried out accordingly to the recommended practice for bread wheat except the different levels of blended NPSB fertilizer rates were studied.

### ***Data collection and measurements***

Days to 50% heading: Days to heading was recorded by counting number of days from the date of sowing until when 50% of the plants in a plot produced spikes above the sheath of the flag leaf that was determined by visual observation.

**Days to 90% physiological maturity:** Days to physiological maturity was recorded by counting the number of days from date of sowing until when 90% of the plants changed green colour to yellowish, loose its water content and attain to physiological maturity in each plot.

**Grain filling period:** Grain filling period was recorded by counting the number of days from heading to maturity, *i.e.* the number of days to maturity minus the number of days to heading.

**Plant height (cm):** It was measured in cm from ground level to the top of the spike excluding the awns from ten randomly pre-tagged plants at maturity time.

**Spike length (cm):** The main spikes from the ten sampled plants were measured in cm and the average represents the spike length in cm for each replication.

**Number of kernels per spike:** Number of kernels per spike was counted from ten randomly sampled plants.

**Grain yield (kg):** The plants in the central rows were harvested and threshed after sun dried. Grain yield per hectare was calculated based on grain yield per plot and it was converted to kg ha<sup>-1</sup>.

**1000 kernels weight (g):** It was determined based on the weight of 1000 seeds sample from the grain yields of each treatment by counting using an electric seed counter and weighted with an electronic balance and the weight was adjusted to 12.5% moisture level.

**Germination test:** The working samples consisted of 300 pure seeds in three replicates of 100 seed each was germinated in double moist filter paper. The seed was placed in each of the petri dishes at a temperature of 25°C under optimum relative humidity for eight days. The seed was checked every day and the papers moistened with distilled water. At the end of the germination period, the seedlings were recorded as normal seedlings; abnormal seedlings; dead seeds and fresh seeds in accordance with the protocol described in [12]. The normal seedlings were considered those possessing well-developed and vigorous shoot and root system. Seeds, which germinated, but did not fall to the normal seedlings, were categorized as abnormal seedlings. Fresh seeds are seeds which remain firm and apparently viable and un-germinated after the prescribed period. Dead seeds are one type of seeds which at the end of test period are neither hard nor fresh and have not produced seedlings. The present germination was determined by considering only the proportion of normal seedlings as follows:

$$\text{Germination (\%)} = \frac{\text{Number of normal seedlings}}{\text{Total number of seeds sown}} \times 100$$

**Speed of germination:** The same procedures were used with that of germination percentage, but the only difference was, the number of normal germinated seeds was recorded daily until there was no further germination. The germinated seedlings in each day were removed after recording. The speed of germination was calculated by adding the quotients of daily counts of normal seedlings divided by the number of day's germination.

$$\text{Speed of Germination} = \frac{N1}{C1} + \frac{N2}{C2} + \dots + \frac{NF}{CF}$$

Where: N1= number of normal seedlings at first count, N2= number of normal seedlings at second count, NF= number of normal seedlings at final count, C1= days to the first count, C2= days to the second count and CF= days to the final count.

**Seedling shoots and root length (cm):** Seeds from the different treatments were germinated as per standard germination test. At the end of the germination period, shoot and root length of five normal seedlings was measured. Shoot length was measured from the point of attachment to the seed to the tip of the seedling, and the average shoot length was calculated by dividing the total shoot length by the number of normal seedlings measured. Similarly, the root length was measured from the point of attachment to the seed to the tip of the root, and the average root

length was computed by dividing the total root length by the total number of the normal seedlings as per [12].

Seedling vigor index i: To determine the seedling vigor index I, seedling lengths (shoot and root length) were measured at the day of final count and determined by using the method developed by [12] as follows: Seed vigor index I = GP x SL

Where: GP = Germination Percentage; SL = Seedling Length

Seedling vigor index ii: To determine the seed vigor index II, seedling dry weight was taken after the final count in grams using a sensitive balance and determined by using the method developed by [12] as follows: Seed vigor index II = GP x SDW

Where: GP = Germination Percentage; SDW = Seedling Dry Weight

Seedling dry weight (g): The weight of five normal seedlings randomly taken from normal germination percentage excluding the cotyledons was oven dried at 80°C for 24 hours and then the seedlings transferred from the oven to desiccators with active silica gel and left for 30 minutes. The seedling dry weight was determined after the final count in grams using a sensitive balance as per the [12] procedure and the average seedling dry weight was calculated.

### **Statistical data analysis**

Data were subjected to analysis of variance (ANOVA) using Gen Stat 15<sup>th</sup> edition statistical software package [13]. All treatment means were compared using the least significant difference (LSD) test at 5% level of significance

## **RESULTS AND DISCUSSION**

### **Soil analysis**

A composite sample was taken from 30 cm depth of the experimental site before sowing of the crop and the physico-chemical properties of the soil were analyzed at Wolaita Sodo soil laboratory. On the basis of particle size distribution, the soil contained 52% sand, 30% silt and 18% clay (Table 1).

**Table 1.** Physico-chemical properties of the top soil (0 to 30 cm) of the experimental fields

Properties	Contents/ Values	Rating
Soil Ph	5.7	Moderately acidic
Organic carbon (%)	1.89	Medium
Total nitrogen (%)	0.169	Low
Available phosphorus (ppm)	7.570	Low
Exchangeable potassium (ppm)	3.153	Very high
Available Sulfur (mg kg <sup>-1</sup> )	17.18	Low
Available Boron (mg kg <sup>-1</sup> )	0.65	Medium
Particle size proportion (%)		
Sand	52	High
Silt	30	Moderate
Clay	18	Low
Soil texture	Loam	

[14] Classified soil organic carbon percentages of <1.0, 1.0 – 1.71, 1.72 – 3.0, 3.1 – 4.29, and >4.3 as very low, low, medium, high, and very high, respectively. Similarly, the authors classified soil total nitrogen percentages of < 0.05, 0.05 – 0.15, 0.15 – 0.25, 0.25 – 0.5, and >0.5

as very low, low, medium, high, and very high, respectively. The authors also classified available phosphorus contents of <5, 5 – 10, 10 – 17, 17 – 25, and >25 mg/kg soil as very low, low, medium, high, and very high, respectively. Similarly, they classified exchangeable potassium contents of <0.2, 0.2 – 0.3, 0.3 – 0.7, 0.7 – 2.0, and > 2.0 as very low, low, medium, high, and very high, respectively. Based on this, the organic carbon at Sadicho district was medium. Available phosphorus and total nitrogen were low, while exchangeable potassium was very high at experimental site. According to the soil textural class determination triangle, the soil of the experimental site is loam which is suitable for bread wheat production [15].

### ***Days to 50% heading, 90% physiological maturity and grain filling period***

Analysis of variance revealed that, days to 50% heading was significantly affected by the main effect of NPSB fertilizer rates, seed sources (SS) and varieties (V). The main effect of NPSB fertilizer rates and seed sources (SS) and their interaction were significant on days to 90% physiological maturity. The main effect of NPSB fertilizer rates and the interaction of NPSB fertilizer and seed sources (NPSB x SS) were highly significant ( $P < 0.01$ ).

The delayed days to 50% heading (58.33 days) and days to 90% physiological maturity (73.00) period were observed in plots that received 125 kg NPSB ha<sup>-1</sup> fertilizer application and seeds obtained from research center, while farmers seed 3 (FC<sub>3</sub>) and control enhanced days to 50% heading (40.56 days) and days to 90% physiological maturity (57.56 days) (Table 2). Delay of treated mean for days to heading and days to maturity might be due to adequate blended NPSB fertilizer has a great contribution to the vigorous vegetative growth and development. The result is in conformity with the findings of [16] who reported that the longest days to heading (71.7) and physiological maturity (113.0) was observed at 300 kg of NPSB application with supplementary urea. The result of this study is also in agreement with the findings of [17] that indicated the significant influence of blended NPS fertilizer on days to heading. The longest duration (24.89 days) of grain filling period was recorded in interaction of plots that received 125 kg NPSB ha<sup>-1</sup> fertilizer and seeds obtained from research center, while the shortest duration (12.33 days) of grain filling period was recorded from farmers seed 2 (FC<sub>2</sub>) and non fertilized once (Table 2). Bread wheat varieties with the longest grain filling period might be due to the time needed for plants to uptake the nutrient in the soil. This result was in agreement with the finding of [18] who reported that increasing the rates of N supplemented from nil to the highest 92 kg N ha<sup>-1</sup> increased days for grain filling period by about 7.76%.

**Table 2.** Interaction effect of NPSB fertilizer and seed sources on days to 50% heading, days to 90% physiological maturity and grain filling period of bread wheat varieties at Sadicho

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources								
	Days to 50% heading			Days to 90% maturity			Grain filling period		
	FC <sub>3</sub>	FC <sub>2</sub>	R	FC <sub>3</sub>	FC <sub>2</sub>	R	FC <sub>3</sub>	FC <sub>2</sub>	R
0	40.56 <sup>a</sup>	42.89 <sup>ab</sup>	40.67 <sup>a</sup>	57.56 <sup>a</sup>	58.89 <sup>ab</sup>	59.56 <sup>abc</sup>	13.00 <sup>a</sup>	12.33 <sup>a</sup>	13.67 <sup>a</sup>
75	42.56 <sup>ab</sup>	41.11 <sup>a</sup>	49.11 <sup>b</sup>	58.89 <sup>ab</sup>	60.44 <sup>abc</sup>	59.00 <sup>ab</sup>	18.11 <sup>bcd</sup>	17.78 <sup>bc</sup>	16.22 <sup>b</sup>
100	45.22 <sup>ab</sup>	47.22 <sup>ab</sup>	46.78 <sup>ab</sup>	62.33 <sup>cd</sup>	62.11 <sup>cd</sup>	64.33 <sup>d</sup>	17.11 <sup>b</sup>	19.44 <sup>cde</sup>	20.56 <sup>e</sup>
125	43.56 <sup>ab</sup>	42.56 <sup>ab</sup>	58.33 <sup>c</sup>	61.11 <sup>bc</sup>	60.11 <sup>abc</sup>	73.00 <sup>e</sup>	20.22 <sup>de</sup>	20.89 <sup>e</sup>	24.89 <sup>f</sup>
LSD (0.05)	5.99			2.74			2.09		
CV (%)	14.2			4.8			12.5		

### Plant height

The analysis of variance for plant height showed significantly affected by the main effect of the NPSB fertilizers and interaction of NPSB fertilizer and seed sources. However, the main effect of seed sources and varieties and all interactions except (NPSB x SS) were non-significant. The highest plant height (154.8 cm) was recorded from the interaction of the highest blended fertilizer rate (125 kg NPSB ha<sup>-1</sup>) and seeds obtained from research center, while the shortest plant height (124.2 cm) was obtained from the control plot (0 kg NPSB ha<sup>-1</sup>) and seeds obtained from farmers seed 2 (FC<sub>2</sub>) (Table 3). The increased plant height in response to increasing rate of blended NPSB fertilizer application from control to 125 kg NPSB ha<sup>-1</sup> was probably due the vital role of nitrogen fertilizer in promoting the vegetative growth and resulted in significant increase in plant height. Seeds obtained from research center play a catalytic role i.e. high genetic purity and identity, as well as high physical, physiological and health quality. This result is in agreement with that of [19] who reported that application of N and P fertilizer rates highly significantly increased plant height of bread wheat. [20] also reported that application of blended fertilizer NPKSBZn (26:11:11:3.5:0.15:0.6) kg ha<sup>-1</sup> with micronutrient Cu+Zn (5+5 ha<sup>-1</sup>) increased plant height of maize by 66.81% over control plot and 6.11% over the recommended NP fertilizers.

### Number of kernels per spike

The main effect of NPSB fertilizer rates and interaction of NPSB fertilizer with seed source (NPSB x SS) and seed source with varieties (SS x V) were significant. Number of kernels per spike is an important yield participating parameter and has a direct consequence on the grain yield of wheat. Higher number of kernels per spike (76.78) was recorded from the interaction of the highest blended fertilizer rate (125 kg NPSB ha<sup>-1</sup>) and seeds obtained from research center, while the lower number of kernels per spike (36.22) was recorded from the control treatment and seeds obtained from farmers seed 2 (FC<sub>2</sub>) (Table 3). The number of kernels per spike was more enhanced by NPSB might be due to the fact that P is essential in development of grains. This result is in agreement with that of [21] who reported that the number of seed per spike becomes significantly increased as the rate of NPSB application increased from 0 to 300 kg ha<sup>-1</sup>. But disagree with that of [18] who reported that non-response of number of kernels per spike to applied NPS and N rates.

**Table 3.** Interaction effect of NPSB fertilizer and seed sources on plant height and number of kernels per spike of bread wheat varieties

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources			Number of kernels per spike		
	Plant height					
	FC <sub>3</sub>	FC <sub>2</sub>	R	FC <sub>3</sub>	FC <sub>2</sub>	R
0	126.1 <sup>a</sup>	124.2 <sup>a</sup>	133.7 <sup>b</sup>	36.78 <sup>a</sup>	36.22 <sup>a</sup>	40.11 <sup>ab</sup>
75	143.5 <sup>cd</sup>	142.7 <sup>cd</sup>	140.7 <sup>c</sup>	57.96 <sup>cd</sup>	48.90 <sup>bc</sup>	53.56 <sup>cd</sup>
100	151.7 <sup>ef</sup>	150.7 <sup>ef</sup>	147.1 <sup>de</sup>	55.69 <sup>cd</sup>	60.31 <sup>d</sup>	59.71 <sup>d</sup>
125	150.0 <sup>ef</sup>	147.4 <sup>de</sup>	154.8 <sup>f</sup>	56.22 <sup>cd</sup>	59.27 <sup>d</sup>	76.78 <sup>e</sup>
LSD (0.05)	5.495			9.074		
CV (%)	4.1			18.1		

### Spike length

Analysis of variance indicated that the main effect of NPSB fertilizer, seed sources and their interaction and 3-way interaction (NPSB x SS x V) were significant. However, main effects of variety and interaction of NPSB fertilizer and variety (NPSB x V) and seed source and variety (SS x V) were non-significant. Interaction of NPSB fertilizer rate, seed source and variety was significant on spike length. The tallest spike length (16 cm) was recorded from the treatment combinations with highest NPSB fertilizer rate (125 kg NPSB ha<sup>-1</sup>), seeds obtained from research center and Hidase variety, while the shortest spike length (5.567 cm) was obtained from control treatment, seed obtained from farmers seed 2 (FC<sub>2</sub>) and Digalu variety (Table 4). This may be due to different varieties may have different yield performing abilities in genetically and increase in spike length at the highest NPSB rate might have resulted from improved root growth and increased uptake of nutrients and better growth favoured due to the synergetic effect of the nutrients at the highest rate. This result agrees with the findings of [22] that indicated NPS fertilizer had ability to increase spike length positively.

**Table 4.** Interaction effect of NPSB fertilizer, seed sources and varieties on spike length of bread wheat varieties

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources	Varieties		
		Digalu	Ogollocho	Hidase
0	FC <sub>3</sub>	5.917 <sup>ab</sup>	6.167 <sup>a-d</sup>	6.500 <sup>a-f</sup>
	FC <sub>2</sub>	5.567 <sup>a</sup>	6.417 <sup>a-f</sup>	6.267 <sup>a-e</sup>
	R	6.167 <sup>a-d</sup>	6.000 <sup>abc</sup>	6.500 <sup>a-f</sup>
75	FC <sub>3</sub>	7.600 <sup>b-g</sup>	7.383 <sup>a-g</sup>	7.700 <sup>b-g</sup>
	FC <sub>2</sub>	7.133 <sup>a-g</sup>	6.900 <sup>a-f</sup>	7.500 <sup>a-g</sup>
	R	7.100 <sup>a-g</sup>	7.617 <sup>b-g</sup>	7.867 <sup>b-g</sup>
100	FC <sub>3</sub>	9.000 <sup>gh</sup>	7.967 <sup>c-g</sup>	7.983 <sup>c-g</sup>
	FC <sub>2</sub>	7.767 <sup>b-g</sup>	8.100 <sup>d-g</sup>	8.200 <sup>efg</sup>
	R	8.350 <sup>fg</sup>	8.217 <sup>efg</sup>	8.033 <sup>d-g</sup>
125	FC <sub>3</sub>	7.833 <sup>b-g</sup>	7.867 <sup>b-g</sup>	7.900 <sup>b-g</sup>
	FC <sub>2</sub>	8.167 <sup>d-g</sup>	8.167 <sup>d-g</sup>	7.967 <sup>c-g</sup>
	R	10.433 <sup>h</sup>	14.833 <sup>i</sup>	16.000 <sup>i</sup>
LSD (0.05)	1.6194			
CV (%)	12.6			

### Grain yield

Analysis of variance indicated that the main effect of NPSB fertilizer, interaction of NPSB fertilizer and seed source (NPSB x SS) and 3-way interaction (NPSB x SS x V) were significant.

The ultimate goal in crop production is maximum economic yield, which is a complex function of individual yield components in response to the genetic potential of the varieties and inputs used. The maximum grain yield (6740.0 kg ha<sup>-1</sup>) was obtained from 3-way interaction in response to application of 125 kg ha<sup>-1</sup> blended NPSB fertilizer, seeds obtained from research center and Hidase variety, while the minimum grain yield (1900.3 kg ha<sup>-1</sup>) was obtained from control treatment, seed obtained from farmer's seed 3 (FC<sub>3</sub>) and Ogollocho variety (Table 5). The minimum grain yield obtained from farmer's seed 3 and Ogollocho variety might be due to low adaptation to agro ecology and poor quality seed sources with respect to agronomic managements. This result agrees with the findings of [23] who reported that using of Kersey Local Seed Business Project seed increases yield by about 23.14% over farmers saved seeds. The highest grain yield at the highest NPSB rates might have resulted from improved root growth and



increased uptake of nutrients and better growth favored due to synergetic effect of the four nutrients which enhanced yield components and yield. Nitrogen affects the vegetative as well as yields whereas phosphorus plays a fundamental role in metabolism and energy producing reaction and can withstand the adverse environmental conditions, thus resulting in enhanced grain yield. The result is in conformity with the findings of [18] who reported that increasing the rates of blended NPS fertilizer from 100 to 200 kg ha<sup>-1</sup> and supplemented N from 0 to 92 kg ha<sup>-1</sup>, the grain yield showed consistent increase. Also [19] reported that application of 100 kg N ha<sup>-1</sup> in combination with 13.6 kg S ha<sup>-1</sup> resulted in highest grain yield of 9.26 tons ha<sup>-1</sup> while the lowest 3.47 tons ha<sup>-1</sup> was recorded from control plot for winter wheat.

### Thousand Kernels weight

Analysis of variance indicated that the main effect of NPSB fertilizer, interaction of NPSB fertilizer and seed source (NPSB x SS) and 3-way interaction (NPSB x SS x V) were significant.

The application of increasing blended fertilizer rates in the experimental area increased the seeds size and the amount of powder of bread wheat varieties. The maximum thousand kernels weight (52.33 g) was recorded from the treatment combinations with highest NPSB fertilizer rate (125 kg NPSB ha<sup>-1</sup>), seeds obtained from research center and Hidase variety, while the minimum thousand kernels weight (29.67 g) was obtained from control treatment, seed obtained from farmers seed 2 (FC<sub>2</sub>) and Hidase variety (Table 5). Higher thousand kernels weight might be due to reflection of improved nutrient use efficiency as a result of increased application of blended fertilizer levels. This result is in line with [24] who reported that applying both micro (especially Zn and B) and macro nutrient and when N -level application increase there is a positive impact on yield component of wheat crop especially on 1000 seed weight.

**Table 5.** Interaction effect of NPSB fertilizer, seed sources and varieties on grain yield and 1000 kernels weight of bread wheat varieties

NPSB (kg ha <sup>-1</sup> )	Seed sources	Varieties					
		Grain yield (kg)			1000 kernels weight (g)		
		Digalu	Ogolicho	Hidase	Digalu	Ogolicho	Hidase
0	FC <sub>3</sub>	2820.0 <sup>bc</sup>	1900.3 <sup>a</sup>	1980.0 <sup>a</sup>	39.53 <sup>c-f</sup>	35.57 <sup>bcd</sup>	32.67 <sup>ab</sup>
	FC <sub>2</sub>	2590.7 <sup>abc</sup>	2290.3 <sup>ab</sup>	2090.3 <sup>ab</sup>	34.57 <sup>abc</sup>	40.10 <sup>c-f</sup>	29.67 <sup>a</sup>
	R	2250.7 <sup>ab</sup>	2620.7 <sup>abc</sup>	3130.3 <sup>cd</sup>	35.67 <sup>bcd</sup>	33.00 <sup>ab</sup>	37.43 <sup>b-e</sup>
75	FC <sub>3</sub>	4400.4 <sup>e-h</sup>	4080.0 <sup>efg</sup>	3600.3 <sup>de</sup>	39.60 <sup>c-f</sup>	39.97 <sup>c-f</sup>	43.83 <sup>f-i</sup>
	FC <sub>2</sub>	4200.7 <sup>efg</sup>	4220.0 <sup>efg</sup>	4630.0 <sup>f-i</sup>	43.43 <sup>f-i</sup>	41.40 <sup>e-h</sup>	39.67 <sup>c-f</sup>
	R	4250.3 <sup>efg</sup>	4740.7 <sup>ghi</sup>	4520.0 <sup>fgh</sup>	39.57 <sup>c-f</sup>	44.10 <sup>f-i</sup>	42.83 <sup>e-i</sup>
100	FC <sub>3</sub>	4310.4 <sup>efg</sup>	4310.9 <sup>efg</sup>	4610.4 <sup>f-i</sup>	44.07 <sup>f-i</sup>	46.00 <sup>ghi</sup>	42.37 <sup>e-h</sup>
	FC <sub>2</sub>	4540.0 <sup>fgh</sup>	4630.0 <sup>f-i</sup>	4740.0 <sup>ghi</sup>	40.57 <sup>d-g</sup>	40.37 <sup>d-g</sup>	43.03 <sup>e-i</sup>
	R	4590.0 <sup>fgh</sup>	4420.0 <sup>e-h</sup>	4660.3 <sup>f-i</sup>	40.93 <sup>d-h</sup>	40.63 <sup>d-g</sup>	43.97 <sup>f-i</sup>
125	FC <sub>3</sub>	5250.7 <sup>hij</sup>	5450.3 <sup>ij</sup>	3850.3 <sup>def</sup>	46.50 <sup>hi</sup>	44.27 <sup>f-i</sup>	43.37 <sup>f-i</sup>
	FC <sub>2</sub>	5560.7 <sup>jk</sup>	4770.7 <sup>ghi</sup>	4500.7 <sup>fgh</sup>	41.93 <sup>e-h</sup>	45.33 <sup>f-i</sup>	44.00 <sup>f-i</sup>
	R	6270.0 <sup>kl</sup>	6500.7 <sup>l</sup>	6740.0 <sup>l</sup>	48.50 <sup>ij</sup>	48.43 <sup>ij</sup>	52.33 <sup>j</sup>
LSD (0.05)	72.35			4.704			
CV (%)	10.7			7.0			

Means followed by the same letter (s) within a column are not significantly different at 5% level of significance, R = Research, FC<sub>2</sub> = Farmers seed 2, FC<sub>3</sub> = Farmers seed 3.

### Germination percentage

The analysis of variance for germination percentage showed significantly affected by the main effect of the NPSB fertilizer, seed sources and their interactions. However, the main effect of varieties and all interactions except (NPSB x SS) were non-significant. The maximum (97.11%) germination percentage was observed due to the interaction of the highest blended fertilizer rate (125 kg NPSB ha<sup>-1</sup>) and seeds obtained from research center, while the minimum germination percentage (81.11%) was recorded from the control treatment and seeds obtained from farmers seed 2 (FC<sub>2</sub>) (Table 6). NPSB fertilizer increases, increase germination of the seed. The possible reason for the maximum germination percentage might be due to N, P, B and S are the main components of the cell that facilitate activity during cell division, which is a sequential event in seed germination. The variety obtained from research center, which had high germination percentage could be considered high quality seeds. This is because the germination percentage is one of the seed quality and vigourisity parameters. This result agrees with the findings of [25] who reported that NP fertilizer increase germination by playing an important role in energy supplying and cell biosynthesis reactions, especially during germination seedling establishment.

**Table 6.** Interaction effect of NPSB fertilizer and seed sources on germination test of bread wheat varieties

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources		
	FC <sub>3</sub>	FC <sub>2</sub>	R
0	83.67 <sup>ab</sup>	81.11 <sup>a</sup>	85.67 <sup>abc</sup>
75	94.22 <sup>fg</sup>	89.44 <sup>c-f</sup>	91.00 <sup>def</sup>
100	92.89 <sup>efg</sup>	89.11 <sup>cde</sup>	92.67 <sup>efg</sup>
125	86.89 <sup>bcd</sup>	86.56 <sup>bcd</sup>	97.11 <sup>g</sup>
LSD (0.05)	4.399		
CV (%)	5.3		

### Speed of germination

Analysis of variance indicated that the main effect of NPSB fertilizer, interaction of NPSB fertilizer and seed source (NPSB x SS) and 3-way interaction (NPSB x SS x V) were significant. The maximum (31.25) speed of germination was observed from 3-way interaction in response to application of 125 kg ha<sup>-1</sup> blended NPSB fertilizer, seeds obtained from research center and Hidase variety, while the minimum speed of germination (14.44) was obtained from control treatment, seed obtained from farmers seed 2 (FC<sub>2</sub>) and Ogolicho variety (Table 7). Wheat varieties with the maximum speed of germination might be due to N, P and S are the main components of the cell that facilitate activity during cell division. The variety obtained from research center, which had high speed of germination could be considered high quality seeds. Variation in speed of germination might be inherent nature of the crop of bread wheat varieties. The varieties, which had high speed of the germination, could be considered high quality seeds. This is because the speed of germination is one of the seed quality and vigourisity parameters. This result is in line with the findings of [25] who reported that NP fertilizer increase speed of germination by playing an important role in energy supplying and cell biosynthesis reactions, especially during germination seedling establishment. [26] also showed that quality seed plays an important role in germination and seedling vigour and ultimately for seed yield.

**Table 7.** Interaction effect of NPSB fertilizer, seed sources and varieties on speed of germination of bread wheat varieties

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources	Varieties		
		Digalu	Ogolicho	Hidase
0	FC <sub>3</sub>	20.49 <sup>c-h</sup>	14.77 <sup>ab</sup>	16.07 <sup>a-e</sup>
	FC <sub>2</sub>	15.45 <sup>a-d</sup>	14.44 <sup>a</sup>	15.26 <sup>abc</sup>
	R	14.49 <sup>a</sup>	16.80 <sup>a-f</sup>	15.12 <sup>abc</sup>
75	FC <sub>3</sub>	22.17 <sup>fgh</sup>	23.66 <sup>gh</sup>	24.12 <sup>gh</sup>
	FC <sub>2</sub>	22.41 <sup>fgh</sup>	22.03 <sup>fgh</sup>	20.94 <sup>dh</sup>
	R	22.53 <sup>fgh</sup>	18.71 <sup>a-g</sup>	20.11 <sup>a-h</sup>
100	FC <sub>3</sub>	22.41 <sup>fgh</sup>	23.52 <sup>gh</sup>	22.05 <sup>fgh</sup>
	FC <sub>2</sub>	23.56 <sup>gh</sup>	23.76 <sup>gh</sup>	21.28 <sup>e-h</sup>
	R	24.64 <sup>gh</sup>	24.00 <sup>gh</sup>	23.56 <sup>gh</sup>
125	FC <sub>3</sub>	23.79 <sup>gh</sup>	23.87 <sup>gh</sup>	19.35 <sup>a-g</sup>
	FC <sub>2</sub>	24.42 <sup>gh</sup>	20.30 <sup>b-h</sup>	24.17 <sup>gh</sup>
	R	22.82 <sup>gh</sup>	26.00 <sup>h</sup>	31.25 <sup>i</sup>
LSD (0.05)	4.798			
CV (%)	13.9			

### Seedling shoots and root length

The main effect of NPSB fertilizer rates and 3-way interaction (NPSB x SS x V) were highly significant ( $P < 0.01$ ).

The tallest seedling length (34.17 cm) was recorded from 3-way interaction of 125 kg ha<sup>-1</sup> blended NPSB fertilizer, seeds obtained from research center and Hidase variety, while the shortest seedling length (17.33 cm) was obtained from control treatment, seed obtained from farmers seed 2 (FC<sub>2</sub>) and Hidase variety (Table 8). Seed sources which produced in different management practices have different potential of germination and potentiality of seedling length. Seeds obtained from research center had heavier seed weight results well-developed shoot and root systems that can withstand any adverse conditions and provide better seedling emergence and seedling establishment in the field. This result was in agreement with the findings of [27] who reported that more seedling length and seedling weight of the heavy seeds might be attributed to large food reserves of the seeds.

### Seedling dry weight

Analysis of variance indicated that the main effect of NPSB fertilizer and seed source and interaction of NPSB fertilizer and seed source (NPSB x SS) and 3-way interaction (NPSB x SS x V) were significant. The interaction of NPSB fertilizer, seed source and variety had significant effect on seedling dry weight. The highest seedling dry weight (0.08200 g) was recorded from 3-way interaction of 125 kg ha<sup>-1</sup> blended NPSB fertilizer, seeds obtained from research center and Hidase variety, while the lowest seedling dry weight (0.02133 g) was obtained from control treatment, seed obtained from research and Hidase variety (Table 8). As increasing NPSB fertilizer level seedling dry weight increases. The possible reason for the highest seedling dry weight might be due to NPSB fertilizer had contributions to shoot and root development. This result was in agreement with the finding of [22] who reported that application of 150 kg NPS rate increased seedling dry weight by 26.29% than non-fertilized.

**Table 8.** Interaction effect of NPSB fertilizer, seed sources and varieties on seedling shoots and root length and seedling dry weight of bread wheat

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources	Varieties					
		Seedling length (cm)			Seedling dry weight (g)		
		Digalu	Ogolicho	Hidase	Digalu	Ogolicho	Hidase
0	FC <sub>3</sub>	18.83 <sup>ab</sup>	21.00 <sup>bc</sup>	21.43 <sup>bc</sup>	0.02600 <sup>ab</sup>	0.03167 <sup>a-d</sup>	0.02300 <sup>a</sup>
	FC <sub>2</sub>	25.60 <sup>def</sup>	21.17 <sup>bc</sup>	17.33 <sup>a</sup>	0.02200 <sup>a</sup>	0.02733 <sup>abc</sup>	0.02567 <sup>ab</sup>
	R	21.83 <sup>bc</sup>	23.50 <sup>cd</sup>	21.67 <sup>bc</sup>	0.02733 <sup>abc</sup>	0.03000 <sup>abc</sup>	0.02133 <sup>a</sup>
75	FC <sub>3</sub>	27.40 <sup>ef</sup>	24.53 <sup>cde</sup>	28.17 <sup>ef</sup>	0.03533 <sup>a-d</sup>	0.03567 <sup>a-d</sup>	0.03367 <sup>a-d</sup>
	FC <sub>2</sub>	27.33 <sup>ef</sup>	27.93 <sup>ef</sup>	26.30 <sup>def</sup>	0.03400 <sup>a-d</sup>	0.03400 <sup>a-d</sup>	0.03500 <sup>a-d</sup>
	R	27.57 <sup>ef</sup>	26.10 <sup>def</sup>	26.73 <sup>def</sup>	0.03533 <sup>a-d</sup>	0.04133 <sup>bcd</sup>	0.03567 <sup>a-d</sup>
100	FC <sub>3</sub>	27.67 <sup>ef</sup>	26.40 <sup>def</sup>	27.03 <sup>def</sup>	0.03267 <sup>a-d</sup>	0.03467 <sup>a-d</sup>	0.03600 <sup>a-d</sup>
	FC <sub>2</sub>	26.73 <sup>def</sup>	26.37 <sup>def</sup>	29.07 <sup>f</sup>	0.03400 <sup>a-d</sup>	0.04467 <sup>cde</sup>	0.04333 <sup>bcd</sup>
	R	27.67 <sup>ef</sup>	28.00 <sup>ef</sup>	28.03 <sup>ef</sup>	0.04167 <sup>bcd</sup>	0.04800 <sup>de</sup>	0.03400 <sup>a-d</sup>
125	FC <sub>3</sub>	28.27 <sup>ef</sup>	29.20 <sup>f</sup>	27.93 <sup>ef</sup>	0.03800 <sup>a-d</sup>	0.03367 <sup>a-d</sup>	0.03300 <sup>a-d</sup>
	FC <sub>2</sub>	28.20 <sup>ef</sup>	27.90 <sup>ef</sup>	26.13 <sup>def</sup>	0.03833 <sup>a-d</sup>	0.03600 <sup>a-d</sup>	0.03000 <sup>abc</sup>
	R	28.50 <sup>f</sup>	27.90 <sup>ef</sup>	34.17 <sup>g</sup>	0.03167 <sup>a-d</sup>	0.05933 <sup>e</sup>	0.08200 <sup>f</sup>
LSD (0.05)	3.140				0.014208		
CV (%)	7.4				24.4		

### Seedling vigor index

Analysis of variance indicated that the main effect of NPSB fertilizer, interaction of NPSB fertilizer and seed source (NPSB x SS) and 3-way interaction (NPSB x SS x V) were significant on seedling vigour index i and ii. The main effect of seed source was also significant on seedling vigour index ii. As increasing NPSB fertilizer level, speed of germination, germination percentage, seedling length and seedling dry weight increases and consequently increase vigourity. Seed obtained from farmers seed 2 (FC<sub>2</sub>) had lower vigor index i and ii as compared to research seed (Table 9). The possible reason for the lowest vigor index i and ii might be due to its lower standard germination percentages, hundred kernel weight, seedling length and seedling dry weight. This result was in conformity with that of [26] who reported that with increase in thousand grain weight, significantly increased seedling length and dry weight that was effective measures of vigor.

**Table 9.** Interaction effect of NPSB fertilizer, seed sources and varieties on seedling vigor index i and ii of bread wheat varieties

NPSB fertilizer (kg ha <sup>-1</sup> )	Seed sources	Varieties					
		Seedling vigor index I			Seedling vigor index II		
		Digalu	Ogolicho	Hidase	Digalu	Ogolicho	Hidase
0	FC <sub>3</sub>	15.99 <sup>ab</sup>	17.19 <sup>abc</sup>	18.06 <sup>bc</sup>	0.02227 <sup>a-d</sup>	0.02243 <sup>a-e</sup>	0.01948 <sup>abc</sup>
	FC <sub>2</sub>	20.65 <sup>cde</sup>	17.70 <sup>abc</sup>	13.70 <sup>a</sup>	0.01764 <sup>a</sup>	0.02649 <sup>a-f</sup>	0.02034 <sup>abc</sup>
	R	19.13 <sup>bcd</sup>	20.30 <sup>cde</sup>	17.96 <sup>bc</sup>	0.02394 <sup>a-f</sup>	0.02598 <sup>a-f</sup>	0.01777 <sup>ab</sup>
75	FC <sub>3</sub>	26.14 <sup>fg</sup>	23.53 <sup>efg</sup>	25.83 <sup>fg</sup>	0.03372 <sup>b-g</sup>	0.03411 <sup>c-g</sup>	0.03082 <sup>a-g</sup>
	FC <sub>2</sub>	25.27 <sup>fg</sup>	25.57 <sup>fg</sup>	22.38 <sup>def</sup>	0.03133 <sup>a-g</sup>	0.03105 <sup>a-g</sup>	0.02958 <sup>a-g</sup>
	R	24.75 <sup>efg</sup>	23.91 <sup>efg</sup>	24.41 <sup>efg</sup>	0.03177 <sup>a-g</sup>	0.03766 <sup>d-g</sup>	0.03267 <sup>a-g</sup>
100	FC <sub>3</sub>	26.17 <sup>fg</sup>	24.41 <sup>efg</sup>	24.70 <sup>efg</sup>	0.03095 <sup>a-g</sup>	0.03215 <sup>a-g</sup>	0.03288 <sup>a-g</sup>
	FC <sub>2</sub>	24.50 <sup>efg</sup>	23.15 <sup>d-g</sup>	25.73 <sup>fg</sup>	0.03105 <sup>a-g</sup>	0.03911 <sup>fg</sup>	0.03837 <sup>efg</sup>
	R	24.18 <sup>efg</sup>	26.49 <sup>fg</sup>	27.01 <sup>fg</sup>	0.03648 <sup>d-g</sup>	0.04459 <sup>g</sup>	0.03275 <sup>a-g</sup>
125	FC <sub>3</sub>	24.78 <sup>efg</sup>	25.48 <sup>fg</sup>	23.90 <sup>efg</sup>	0.03329 <sup>a-g</sup>	0.02940 <sup>a-g</sup>	0.02831 <sup>a-f</sup>
	FC <sub>2</sub>	24.78 <sup>efg</sup>	23.96 <sup>efg</sup>	22.45 <sup>def</sup>	0.03371 <sup>b-g</sup>	0.03096 <sup>a-g</sup>	0.02571 <sup>a-f</sup>
	R	26.96 <sup>fg</sup>	27.26 <sup>g</sup>	33.83 <sup>h</sup>	0.02990 <sup>a-g</sup>	0.05811 <sup>h</sup>	0.08118 <sup>i</sup>
LSD (0.05)	3.754				0.012845		
CV (%)	9.9				24.5		

Means followed by the same letter (s) within a table are not significantly different at 5% level of significance, R = Research, FC<sub>2</sub> = Farmers seed 2, FC<sub>3</sub> = Farmers seed 3

## CONCLUSIONS

Inappropriate uses of fertilizer rate, unavailability of quality seed for varieties that are high yielding as well as adapted to wide range of agro-ecologies of the area and the use of farm saved poor quality seeds of local varieties are some of the major factors which limit yield of wheat. The results of yield components and seed quality investigations showed that days to 50% heading, maturity, grain filling period, plant height, number of kernels per spike and germination percentage were significantly affected by the interaction effects of fertilizer rate and seed sources, while spike length, grain yield, 1000 kernels weight, speed of germination, seedling dry weight, seedling length, seedling vigor index i and ii were significantly affected by 3-way interaction. The maximum values of all parameters recorded in interaction of 125 kg ha<sup>-1</sup> blended NPSB fertilizer, seeds obtained from research center and Hidase variety. Therefore, application of NPSB at the rate of 125 kg ha<sup>-1</sup>, seeds obtained from research center and Hidase variety were economically profitable and recommended for farmers around Sadicho district and other areas with similar agro-ecology.

**Acknowledgment.** The authors are grateful to Mr. Liranso Habdolo and W/Kidan Maryo for their considerable support in trials management and data collection.

## REFERENCES

- [1] Jemal, A., Tamado, T. and Firdissa, E. 2015: Response of Bread Wheat (*Triticum aestivum* L.) Varieties to Seeding Rates at Kulumsa, Southeastern Ethiopia. *Asian Journal of Plant Sciences*, 14: 5058.
- [2] FAOSTAT (Food and Agricultural Organization of the United Nations Statistics), (2015): Food balance sheets.
- [3] CIMMYT (International Maize and Wheat Improvement Centre). 2011: Wheat Global alliance for improving food security and the livelihoods of the resource poor in the developing world. Proposal submitted by CIMMYT and ICARDA to the CGIAR Consortium Board. August 30, 2011.
- [4] Central Statistical Agency of Ethiopia (CSA), 2019: Agricultural sample survey Report on area and production for major crops 1: 584.
- [5] Bewley, JD. and Black, M. 1994: *Seeds Physiology of Development and Germination, Second edition*. Plenum Press, New York. Ali, R., M.J. Khan and R.A. Khattak. 2008: Response of rice to different sources of Sulfur (S) at various levels and its residual effect on wheat in rice-wheat cropping system. *Soil Environment* 27(1): 131-137.
- [6] Haile, W., & Boke, S. (2011): Response of Irish potato (*Solanum tuberosum*) to the application of potassium at acidic soils of chench, Southern Ethiopia. *International Journal of Agricultural Biology*, 13, 595–598.
- [7] EthioSIS (Ethiopia Soil Information System). 2014: Soil fertility status and fertilizer recommendation atlas for Southern regional state, Ethiopia.
- [8] Dawit, H., Kasu, T., Wubengeda, A., Tadese, D. and Asrat M. 2015: Agronomic and Economic Evaluation of the N and P Response of Bread Wheat Grown in Moist and Humid Mid highland Vertisols Areas of Arsi Zone, Ethiopia. *African Journal of Agricultural Research*, 10 (3):89-9946
- [9] Ziadi, N., G., Belander, A. N., Cambouris, N., Tremblay, M. C., Nolinand A., Claessens. 2008: Relationship between phosphorus and nitrogen concentrations in spring wheat. *Agronomy Journal*, 100(1):80-86.
- [10] Marschner, H., 1995: Mineral nutrition of higher plants. 2nd Ed. New York: Academic Press, pp: 889.

- [11] Doyogana Woreda Agriculture and Natural Resource Office report (2020) :Annual report, Doyogana, Ethiopia.
- [12] ISTA (International Rules for Seed Testing Association).2014: *International Seed Testing Association*, Bassersdorf, Switzerland
- [13] GenStat. 2012: GenStat Procedure Library Release. 15<sup>th</sup> edition. VSN International Ltd.
- [14] Hazelton, P., and Murphy, B. 2007: Interpreting soil test results: What do all the numbers mean? Second Edition. CSIRO Publishing, Pp.152.
- [15] Miyan MS., Impiglia A., Anderson WK. (2011): Agronomic practices for durum wheat in an area new to the crop. *Communications in Biometry and Crop Science* 6(2):64-79.
- [16] Diriba, S., G., Rut-Duga, D. and Wogayehu, W. 2018: Effects of Blended Fertilizer Rates on Bread Wheat (*Triticum Aestivum* L.) Varieties on Growth and Yield Attributes. *Journal of Ecology and Natural Resources*, 3(3): 000170
- [17] Tagesse A., Ketema B and Tamado T. 2017: Effect of Blended NPS Fertilizer Supplemented with Nitrogen on Yield Components and Yield of Bread Wheat (*Triticum aestivum* L.). *Journal of Natural Sciences Research*, 8(11):90-96
- [18] Tilahun, A and Tamado, T. 2016: Growth, yield component and yield response of durum wheat (*Triticum turgidum* L. var. *Durum*) to blended NPS fertilizer supplemented with N rates at Arsi Negelle, Central Ethiopia. *African Journal of Plant Science* 13(1):9 -20
- [19] Melesse, H. (2017): Response of bread wheat (*Triticum aestivum* L.) varieties to N and P fertilizer rates in Ofla district, Southern Tigray, Ethiopia. *African Journal of Agricultural Research* 12(19):1646-1660.
- [20] Dagne, C. (2016): Blended Fertilizers Effects on Maize Yield and Yield Components of Western Oromia, Ethiopia. *Agriculture, Forestry and Fisheries* 5(5): 151-162.
- [21] Rut-Duga D., Diriba Sh. and Wogayehu W. 2019: Effects of Blended Fertilizer Rates on Bread Wheat (*Triticum aestivum* L.) Varieties on Growth and Yield Attributes *Journal of Ecology and Natural Resources*. 3(3): 000170
- [22] Demisew, A. (2017): Effect of Blended NPS Fertilizer on Seed Quality, Yield and Yield Related Traits of Bread Wheat (*Triticum aestivum* L.) Varieties at Haramaya, MSc Thesis, Haramaya University.
- [23] Alemayehu, A. (2015): Effect of Seed Sources and Rates on Productivity of Bread Wheat (*Triticum aestivum* L.) Varieties at Kersa, Eastern Ethiopia, MSc Thesis, Haramaya University.
- [24] Muhammad T., Asefa, T., Tajamol, H., Wasoya, A. (2009): Yield response of wheat to Boron application. *Pak J Life Soc Sci* 7(1): 39-42.
- [25] White, P.J., Veneklaas, E.J. 2012: Nature and nurture: the importance of seed phosphorus content. *Plant Soil*, 357:1-8.
- [26] Khan, ML. 2003: Effects of seed mass on seedling success in *Artocarpus heterophyllus* L. a tropical tree species of north- east India. *Indian Journal of Agricultural Science*, 25: 103-110.
- [27] Gharineh., M.H. and Moshatati, A. 2012: Effect of grain weight on germination and seed vigour of wheat. *International Journal of Agriculture and Crop Sciences*, 4 (8): 458-460.